

Exam 3 Fall 24 Answer Key

$$1(a) \sum_{n=1}^{\infty} \frac{(-1)^n (3x+5)^n}{(3n+5)^2 \cdot 4^n}$$

Ratio Test

Converges by Ratio Test when

$$L = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} (3x+5)^{n+1}}{(3(n+1)+5)^2 \cdot 4^{n+1}} \cdot \frac{(3n+5)^2 \cdot 4^n}{(-1)^n (3x+5)^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(3x+5)^{n+1}}{(3x+5)^n} \cdot \left(\frac{3n+5}{3n+8} \right)^2 \cdot \frac{4^n}{4^{n+1}} \right| = \frac{|3x+5|}{4} < 1$$

$$\frac{|3x+5|}{4} < 1 \Rightarrow |3x+5| < 4 \Rightarrow -4 < 3x+5 < 4 \Rightarrow -9 < 3x < -1 \Rightarrow -3 < x < -\frac{1}{3}$$

Manually Check Convergence at Endpoints

Take $x = -3$ Series becomes

$$\sum_{n=1}^{\infty} \frac{(-1)^n [3(-3)+5]^n}{(3n+5)^2 \cdot 4^n} = \sum_{n=1}^{\infty} \frac{(-1)^n (-4)^n}{(3n+5)^2 \cdot 4^n} = \sum_{n=1}^{\infty} \frac{(-1)^n (-1)^n 4^n}{(3n+5)^2 \cdot 4^n} = \sum_{n=1}^{\infty} \frac{1}{(3n+5)^2} \approx \sum_{n=1}^{\infty} \frac{1}{n^2}$$

Convergent p-Series $p = 2 > 1$

Bound Terms

$$\frac{1}{(3n+5)^2} \leq \frac{1}{n^2}$$

\Rightarrow Series Converges by CT

OR LCT Limit

$$\lim_{n \rightarrow \infty} \frac{(3n+5)^2}{1/n^2} = \lim_{n \rightarrow \infty} \frac{n^2}{(3n+5)^2} = \lim_{n \rightarrow \infty} \left(\frac{n}{3n+5} \right)^2 = \frac{1}{9} \text{ Finite Non-zero}$$

\Rightarrow Series Converges by LCT

Take $x = -\frac{1}{3}$ Series becomes

$$\sum_{n=1}^{\infty} \frac{(-1)^n [3(-\frac{1}{3})+5]^n}{(3n+5)^2 \cdot 4^n} = \sum_{n=1}^{\infty} \frac{(-1)^n 4^n}{(3n+5)^2 \cdot 4^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{(3n+5)^2} \xrightarrow{AS} \sum_{n=1}^{\infty} \frac{1}{(3n+5)^2}$$

Already Shown above

to be Convergent using CT or LCT

ACT

Original Series

Converges by ACT

OR

$$1. \text{ Isolate } b_n = \frac{1}{(3n+5)^2} > 0$$

$$2. \lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \frac{1}{(3n+5)^2} = 0$$

3. Terms Decreasing


$$b_{n+1} = \frac{1}{(3(n+1)+5)^2} = \frac{1}{(3n+8)^2} \leq \frac{1}{(3n+5)^2} = b_n$$

Series Converges by AST

Finally, Interval of Convergence
Radius of Convergence

$$I = \left[-3, -\frac{1}{3} \right]$$

$$R = \frac{4}{3}$$



1(b) Multiple choices

$$\sum_{n=1}^{\infty} n^n (x-6)^n$$

Ratio Test

$$L = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{\overset{(n+1)^n(n+1)}{(n+1)^{n+1}} (x-6)^{n+1}}{n^n (x-6)^n} \right| = \lim_{n \rightarrow \infty} \frac{\overset{e}{(n+1)^n}}{n^n} \overset{\infty}{(n+1)} |x-6| = \infty > 1$$

Diverges by Ratio Test
unless $x=6$ (when $L=0<1$)

$$\Rightarrow I = \{6\}$$

$\Rightarrow R=0$ not needed this time

OR others will work ..

$$\sum_{n=1}^{\infty} (2n)! (x-6)^n$$

$$\sum_{n=1}^{\infty} (n!)^2 (x-6)^n$$

$$\sum_{n=1}^{\infty} n! (x-6)^n$$

$$\sum_{n=1}^{\infty} (3n)! n^n (x-6)^n$$

⋮

justify

$$2(a) \ln(1+9x^2) = \sum_{n=0}^{\infty} \frac{(-1)^n (9x^2)^{n+1}}{n+1} = \sum_{n=0}^{\infty} \frac{(-1)^n 9^{n+1} x^{2n+2}}{n+1}$$

$$\text{Need } |9x^2| < 1 \Rightarrow |x|^2 < \frac{1}{9} \Rightarrow |x| < \frac{1}{3} \Rightarrow R = \frac{1}{3}$$

Recall:

$$\ln(1+x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{n+1}}{n+1}$$

$$2(b) \quad x^3 e^{-4x} = x^3 \sum_{n=0}^{\infty} \frac{(-4x)^n}{n!} = x^3 \sum_{n=0}^{\infty} \frac{(-1)^n 4^n x^n}{n!} = \sum_{n=0}^{\infty} \frac{(-1)^n 4^n x^{n+3}}{n!}$$

$R = \infty$ (under x^3)
 $R = \infty$ (under \sum)

$R = \infty$ (circled)

Recall:

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$2(c) \quad \frac{d}{dx} \left(8x^4 \sin(8x) \right) = \frac{d}{dx} \left(8x^4 \sum_{n=0}^{\infty} \frac{(-1)^n (8x)^{2n+1}}{(2n+1)!} \right) = \frac{d}{dx} \left(8x^4 \sum_{n=0}^{\infty} \frac{(-1)^n 8^{2n+1} x^{2n+1}}{(2n+1)!} \right)$$

$$= \frac{d}{dx} \sum_{n=0}^{\infty} \frac{(-1)^n 8^{2n+2} x^{2n+5}}{(2n+1)!} = \sum_{n=0}^{\infty} \frac{(-1)^n 8^{2n+2} (2n+5) x^{2n+4}}{(2n+1)!}$$

$R = \infty$

STILL After Differentiation

Recall:

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$$

$$2(d) \quad \int \frac{x^2}{8+x^3} dx = \int x^2 \left(\frac{1}{8+x^3} \right) dx = \int \frac{x^2}{8} \left(\frac{1}{1 + \frac{x^3}{8}} \right) dx = \int \frac{x^2}{8} \left(\frac{1}{1 - (-\frac{x^3}{8})} \right) dx$$

$$= \int \frac{x^2}{8} \sum_{n=0}^{\infty} \left(\frac{-x^3}{8} \right)^n dx = \int \frac{x^2}{8} \sum_{n=0}^{\infty} \frac{(-1)^n x^{3n}}{8^n} dx = \int \sum_{n=0}^{\infty} \frac{(-1)^n x^{3n+2}}{8^{n+1}} dx$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n x^{3n+3}}{8^{n+1} (3n+3)} + C$$

$$\text{Need } \left| \frac{-x^3}{8} \right| < 1 \Rightarrow |x|^3 < 8$$

$$\Rightarrow |x| < 2$$

$$\text{Recall: } \frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$$

$R = 2$

STILL After Integration

3. $\frac{1}{\sqrt{e}} = e^{-\frac{1}{2}}$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$$

$$16 \cdot 24 = 384$$

$$e^{-1/2} = 1 + \left(-\frac{1}{2}\right) + \frac{\left(-\frac{1}{2}\right)^2}{2!} + \frac{\left(-\frac{1}{2}\right)^3}{3!} + \frac{\left(-\frac{1}{2}\right)^4}{4!} + \dots$$

$$= 1 - \frac{1}{2} + \frac{1}{8} - \frac{1}{48} + \frac{1}{384} - \dots$$

$$\approx 1 - \frac{1}{2} + \frac{1}{8} - \frac{1}{48} = \frac{48 - 24 + 6 - 1}{48} = \frac{29}{48} \leftarrow \text{Estimate}$$

Make
Sure to
Connect
Solution
Throughout

Using the Alternating Series Estimation Theorem (A.S.E.T)

we can Estimate the Full Sum using only the

first four terms with error at most $\frac{1}{384} < \frac{1}{100}$ as desired.

4 a. $\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n}}{3^{2n} (2n+1)!} = \sum_{n=0}^{\infty} \frac{(-1)^n \left(\frac{\pi}{3}\right)^{2n} \cdot \frac{\pi}{3}}{(2n+1)! \cdot \frac{\pi}{3}} = \frac{3}{\pi} \sum_{n=0}^{\infty} \frac{(-1)^n \left(\frac{\pi}{3}\right)^{2n+1}}{(2n+1)!}$

Flip

Recall:

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$$

$$= \frac{3}{\pi} \sin\left(\frac{\pi}{3}\right) = \frac{3\sqrt{3}}{2\pi}$$

4 b. $-1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} - \frac{1}{5} + \frac{1}{6} - \dots = -\left(1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \dots\right)$

$$= -\ln(1+1) = -\ln 2$$

Recall:

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

warning: $\ln(1+(-1)) = \ln 0$ undefined
won't work

4 c. $\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n+1}}{4^n (2n)!} = \pi \sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n}}{2^{2n} (2n)!} = \pi \sum_{n=0}^{\infty} \frac{(-1)^n \left(\frac{\pi}{2}\right)^{2n}}{(2n)!} = \pi \cos\left(\frac{\pi}{2}\right) = \pi \cdot 0 = 0$

Recall:

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$$

Don't Drop

$$4d \sum_{n=0}^{\infty} \frac{(-1)^{n+1} (\ln 9)^n}{2^n \cdot n!} = - \sum_{n=0}^{\infty} \frac{(-1)^n (\ln 9)^n}{2^n n!} = - \sum_{n=0}^{\infty} \frac{\left(\frac{-\ln 9}{2}\right)^n}{n!} = -e^{\frac{-\ln 9}{2}}$$

Recall:

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$= -e^{-\frac{1}{2} \ln 9} = -e^{\ln(9^{-\frac{1}{2}})} = -\frac{1}{9^{\frac{1}{2}}} = -\frac{1}{3}$$

Extra 1

$$4e. 4 + 4 - \frac{4}{3} + \frac{4}{5} - \frac{4}{7} + \frac{4}{9} - \dots = 4 \left(1 + 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots \right)$$

Recall:

$$\arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

$$\arctan 1 = 1 - \frac{1}{3} + \frac{1}{5} - \dots = 4 \left[1 + \arctan(1) \right] = 4 \left(1 + \frac{\pi}{4} \right) = 4 + \pi$$

$$4f. -\frac{\pi^2}{2!} + \frac{\pi^4}{4!} - \frac{\pi^6}{6!} + \dots = (\cos \pi) - 1 = -1 - 1 = -2$$

Recall:

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \quad \longrightarrow \quad \cos \pi = 1 - \frac{\pi^2}{2!} + \frac{\pi^4}{4!} - \dots$$

missing

$$5. \lim_{x \rightarrow 0} \frac{1 - \cos(2x)}{e^{-x} - 1 + x} = \lim_{x \rightarrow 0} \frac{1 - \left(1 - \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} - \frac{(2x)^6}{6!} + \dots \right)}{1 + (-x) + \frac{(-x)^2}{2!} + \frac{(-x)^3}{3!} + \frac{(-x)^4}{4!} + \dots - 1 + x}$$

$$= \lim_{x \rightarrow 0} \frac{1 - 1 + \frac{2^2 x^2}{2!} - \frac{2^4 x^4}{4!} + \frac{2^6 x^6}{6!} - \dots}{1 - x + \frac{x^2}{2!} - \frac{x^3}{3!} + \frac{x^4}{4!} - \dots - 1 + x}$$

Need all ...

$$= \lim_{x \rightarrow 0} \frac{\frac{2^2 x^2}{2!} - \frac{2^4 x^4}{4!} + \frac{2^6 x^6}{6!} - \dots}{\frac{x^2}{2} - \frac{x^3}{3!} + \frac{x^4}{4!} - \dots}$$

1/x^2
1/x^2

Show All Algebra Steps

$$= \lim_{x \rightarrow 0} \frac{\frac{4}{2} - \frac{2^4 x^2}{4!} + \frac{2^6 x^4}{6!} - \dots}{\frac{1}{2} - \frac{x}{3!} + \frac{x^2}{4!} - \dots} = \frac{\frac{4}{2}}{\frac{1}{2}} = 4$$

5. continued



Check answer with Optional L'Hôpital's Rule

$$\lim_{x \rightarrow 0} \frac{1 - \cos(2x)}{e^{-x} - 1 + x} \stackrel{\frac{0}{0}}{=} \lim_{x \rightarrow 0} \frac{2 \sin(2x)}{-e^{-x} + 1} \stackrel{\frac{0}{0}}{=} \lim_{x \rightarrow 0} \frac{4 \cos(2x)}{-e^{-x}} = \frac{4}{1} = 4 \text{ Match!}$$

$$6. \arctan(x^4) = \int \frac{4x^3}{1+x^8} dx = \int 4x^3 \left(\frac{1}{1+x^8} \right) dx \quad \star \text{ Don't Drop } dx's$$

$$= \int 4x^3 \left(\frac{1}{1 - (-x^8)} \right) dx = \int 4x^3 \sum_{n=0}^{\infty} (-x^8)^n dx$$

$$= \int 4x^3 \sum_{n=0}^{\infty} (-1)^n x^{8n} dx = \int 4 \sum_{n=0}^{\infty} (-1)^n x^{8n+3} dx$$

$$= 4 \sum_{n=0}^{\infty} \frac{(-1)^n x^{8n+4}}{8n+4} + C = \sum_{n=0}^{\infty} \frac{(-1)^n x^{8n+4}}{2n+1} + C$$

$\cancel{4} / (2n+1)$

Expand in Long Form to Solve for + C

$$\arctan(x^4) = \frac{x^4}{1} - \frac{x^{12}}{3} + \frac{x^{20}}{5} - \dots + C$$

Test the Center $x=0$ into both sides to solve for + C

$$\arctan 0 = 0 - 0 + 0 - \dots + C \Rightarrow C = 0$$

Finally,

$\arctan(x^4) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{8n+4}}{2n+1}$

 $\stackrel{or}{=} 4 \sum_{n=0}^{\infty} \frac{(-1)^n x^{8n+4}}{8n+4}$

Optional Check using "Substitution" into $\arctan x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1}$

$$\arctan(x^4) = \sum_{n=0}^{\infty} \frac{(-1)^n (x^4)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} \frac{(-1)^n x^{8n+4}}{2n+1}$$

Match!