- This is a closed-book examination. No books, notes, calculators, cell phones, communication devices of any sort, or other aids are permitted.
- You need not simplify algebraically complicated answers. However, numerical answers such as $\sin\left(\frac{\pi}{6}\right)$, $4^{\frac{3}{2}}$, $e^{\ln 4}$, $\ln(e^7)$, $e^{3\ln 3}$, or $\cosh(\ln 3)$ should be simplified.
- ullet Please show all of your work and justify all of your answers. (You may use the backs of pages for additional work space.)
- **1.** [15 Points] Find the Interval and Radius of Convergence for the following power series. Analyze carefully and with full justification.

$$\sum_{n=1}^{\infty} \frac{(-1)^n (5x-2)^n}{(n+5) \ 8^n}$$

2. [12 Points] Find the MacLaurin series representation for each of the following functions. State the Radius of Convergence for each series. Your answer should be in sigma notation $\sum_{i=1}^{n}$

(a)
$$f(x) = \frac{x}{1 + 7x}$$

(b)
$$f(x) = x^4 \arctan(4x)$$

- **3.** [15 Points]
- (a) Write the MacLaurin Series representation for $f(x) = x^3 \ln(1 + x^3)$.
- (b) Use the MacLaurin Series representation for $f(x) = x^3 \ln(1+x^3)$ from part (a) to

Estimate
$$\int_0^1 x^3 \ln(1+x^3) dx$$
 with error less than $\frac{1}{30}$.

Justify in words that your error is indeed less than $\frac{1}{30}$

4. [18 Points] Find the **sum** for each of the following series.

(a)
$$\frac{1}{\pi} - \frac{1}{2(\pi)^2} + \frac{1}{3(\pi)^3} - \frac{1}{4(\pi)^4} + \frac{1}{5(\pi)^5} - \frac{1}{6(\pi)^6} + \dots$$
 (b) $\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n}}{(36)^n (2n+1)!}$

(b)
$$\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n}}{(36)^n (2n+1)!}$$

(c)
$$\sum_{n=0}^{\infty} \frac{(-1)^n (\ln 8)^n}{3^{n+1} n!}$$

(c)
$$\sum_{n=0}^{\infty} \frac{(-1)^n (\ln 8)^n}{3^{n+1} n!}$$
 (d)
$$\frac{1}{\sqrt{3}} - \frac{1}{3(\sqrt{3})^3} + \frac{1}{5(\sqrt{3})^5} - \frac{1}{7(\sqrt{3})^7} + \dots$$

1

(e)
$$\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n}}{2^{2n} (2n)!}$$
 (f) $\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n}}{2^{4n} (2n)!}$

(f)
$$\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n}}{2^{4n} (2n)!}$$

- **5.** [20 Points] Volumes of Revolution
- (a) Consider the region bounded by $y = \arctan x$, y = 8 x, x = 0, and x = 1. Rotate this region about the horizontal line y = -1. Set-up, **BUT DO NOT EVALUATE!!**, the integral to compute the volume of the resulting solid using the Washer Method. Sketch the solid, along with one of the approximating washers.
- (b) Consider the region bounded by $y = 1 + e^x$, y = 4, and x = 0. Rotate this region about the vertical line x = -2. Set-up, **BUT DO NOT EVALUATE!!**, the integral to compute the volume of the resulting solid using the Cylindrical Shells Method. Sketch the solid, along with one of the approximating shells.
- (c) Consider the region bounded by $y = \sin x$, $y = \cos x$, x = 0 and $x = \frac{\pi}{4}$. Rotate this region about the vertical line x = 3. Set-up, **BUT DO NOT EVALUATE!!**, the integral to compute the volume of the resulting solid using the Cylindrical Shells Method. Sketch the solid, along with one of the approximating shells.
- (d) Consider the region bounded by $y = \ln x$, $y = e^x$, x = 1 and x = 3. Rotate this region about the y-axis. COMPUTE the volume of the resulting solid using the Cylindrical Shells Method. Sketch the solid, along with one of the approximating shells.
- 6. [20 Points] Parametric Equations
- (a) Consider the Parametric Curve given by $x = \frac{t^3}{3} \frac{e^{2t}}{2}$ and $y = 2te^t 2e^t$.

<u>COMPUTE</u> the **arclength** of this parametric curve for $0 \le t \le 1$.

(b) Consider a different Parametric Curve given by $x = \sin^3 t$ and $y = \cos^3 t$.

<u>COMPUTE</u> the **surface area** obtained by rotating this curve about the x-axis, for $0 \le t \le \frac{\pi}{2}$.

OPTIONAL BONUS

OPTIONAL BONUS #1 Consider the region bounded by $y = \ln x$, $y = e^x$, x = 1, and x = e. Rotate this region about the y-axis. Compute the resulting volume using two methods: Cylindrical Shells Method and Washer Method.