

Math 121 Final Exam December 19, 2019

- This is a closed-book examination. No books, notes, calculators, cell phones, communication devices of any sort, or other aids are permitted.
- Numerical answers such as $\sin\left(\frac{\pi}{6}\right)$, $4^{\frac{3}{2}}$, $e^{\ln 4}$, $\ln(e^7)$, $e^{-\ln 5}$, $e^{3\ln 3}$, $\arctan(\sqrt{3})$, or $\cosh(\ln 3)$ should be simplified.
- Please *show* all of your work and *justify* all of your answers. (You may use the backs of pages for additional work space.)

1. [12 Points] Evaluate the following **limits**. Please justify your answer. Be clear if the limit equals a value, $+\infty$ or $-\infty$, or Does Not Exist. Simplify.

(a) $\lim_{x \rightarrow 0} \frac{xe^x - \sin x}{\ln(1+x) - \arctan x}$ (b) Compute $\lim_{x \rightarrow 0} \frac{xe^x - \sin x}{\ln(1+x) - \arctan x}$ **again** using series.

2. [18 Points] Evaluate the following **integral**.

(a) Show that $\int_0^{\frac{\pi}{2}} \frac{\cos x}{(1 + \sin^2 x)^{\frac{7}{2}}} dx = \frac{43}{60\sqrt{2}}$ (b) $\int_1^{\sqrt{3}} \frac{x^2}{\sqrt{4-x^2}} dx$

3. [40 Points] For each of the following **improper integrals**, determine whether it converges or diverges. If it converges, find its value. Simplify.

(a) $\int_0^5 \frac{6}{x^2 - 4x - 5} dx$ (b) $\int_0^{e^5} \frac{1}{x [25 + (\ln x)^2]} dx$
 (c) $\int_{-\infty}^5 \frac{6}{x^2 - 4x + 7} dx$ (d) $\int_1^2 \frac{1}{x \ln x} dx$ (e) $\int_0^e \frac{\ln x}{\sqrt{x}} dx$

4. [18 Points] Find the **sum** of each of the following series (which do converge). Simplify.

(a) $\sum_{n=1}^{\infty} \frac{(-3)^n - 2}{4^n}$ (Hint: split?) (b) $\sum_{n=0}^{\infty} \frac{(-1)^{n+1} (\ln 9)^n}{2^{n+1} \cdot n!}$ (c) $\sum_{n=0}^{\infty} \frac{(-1)^{n+1} \pi^{2n-1}}{9^n (2n)!}$
 (d) $\frac{\pi^3}{3!} - \frac{\pi^5}{5!} + \frac{\pi^7}{7!} - \frac{\pi^9}{9!} + \dots$ (e) $-1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} - \frac{1}{5} + \dots$ (f) $3 - 1 + \frac{3}{5} - \frac{3}{7} + \frac{3}{9} - \dots$

5. [24 Points] In each case determine whether the given series is **absolutely convergent**, **conditionally convergent**, or **divergent**. Justify your answers.

(a) $\sum_{n=1}^{\infty} (-1)^n \frac{n^2 + 7}{n^7 + 2}$ (b) $\sum_{n=1}^{\infty} \frac{\arctan n}{7} + \frac{7}{\arctan n}$
 (c) $\sum_{n=1}^{\infty} (-1)^n \left(\frac{\sqrt{n} + 7}{n} \right)$ (d) $\sum_{n=1}^{\infty} \frac{(-1)^n (3n)! \ln n}{(n!)^2 e^{4n} n^n}$

6. [20 Points] Find the **Interval** and **Radius** of Convergence for the following power series. Analyze carefully and with full justification.

$$(a) \sum_{n=1}^{\infty} \frac{(-1)^n (3x - 5)^n}{(n + 7)^2 \cdot 7^{n+1}} \quad (b) \sum_{n=1}^{\infty} \frac{x^{2n+1}}{n^n} \quad (c) \sum_{n=1}^{\infty} n! (x - 6)^n$$

7. [10 Points] Please analyze with detail and justify carefully. Simplify.

(a) Use MacLaurin Series to **Estimate** $\frac{1}{\sqrt{e}}$ with error less than $\frac{1}{100}$.

(b) Compute the MacLaurin Series for $f(x) = \frac{1}{(1-x)^2}$ and then **State** the Radius of Convergence. Your answer should be in Sigma notation. Hint: Use Differentiation.

8. [10 Points] For both parts, you do **not** need to find the Radius of Convergence. Your answer should be in Sigma notation or write out the first 5 non-zero terms.

(a) Demonstrate one method to compute the MacLaurin Series for $f(x) = \ln(1+x)$. Justify. Do not just write down the formula.

(b) Demonstrate a second, **different** method to compute the MacLaurin Series for $f(x) = \ln(1+x)$. Justify. Do not just write down a formula.

9. [10 Points]

(a) Write the **first 6 non-zero terms** of the MacLaurin Series for $f(x) = \sin(x^3) + \cos(x^3)$.

(b) Use this series to now determine the **sixth, seventh, eighth** and **ninth** derivatives of $f(x) = \sin(x^3) + \cos(x^3)$ evaluated at $x = 0$. Do Not Simplify your answers.

10. [18 Points]

(a) Consider the Parametric Curve represented by $x = (\arctan t) - t$ and $y = 2 \sinh^{-1} t$.

Recall $\frac{d}{dx} \sinh^{-1} x = \frac{1}{\sqrt{1+x^2}}$

COMPUTE the **arclength** of this parametric curve for $0 \leq t \leq \sqrt{3}$.

(b) Consider a different Parametric Curve represented by $x = \cos^3 t$ and $y = \sin^3 t$.

COMPUTE the **surface area** obtained by rotating this curve about the y -axis for $0 \leq t \leq \frac{\pi}{2}$.

11. [20 Points] For **each** of the following problems, do the following **THREE** things:

- Sketch the Polar curve(s) and shade the described bounded region.
- Set-Up but **DO NOT EVALUATE** an Integral representing the area of the described bounded region.
- Set-Up but **DO NOT EVALUATE** another **slightly different** Integral representing the same area of the described bounded region.
 - The **area** bounded outside the polar curve $r = 3 + 3 \cos \theta$ and inside $r = 9 \cos \theta$.
 - The **area** bounded outside the polar curve $r = 1$ and inside the polar curve $r = 2 \sin \theta$.
 - The **area** that lies inside both of the curves $r = 2 + 2 \sin \theta$ and $r = 2 - 2 \sin \theta$.
 - The **area** bounded inside **one** petal of the curve $r = 3 \sin(2\theta)$.