Math 121 Final Exam May 7, 2018

- This is a closed-book examination. No books, notes, calculators, cell phones, communication devices of any sort, or other aids are permitted.
- Numerical answers such as $\sin\left(\frac{\pi}{6}\right)$, $4^{\frac{3}{2}}$, $e^{\ln 4}$, $\ln(e^7)$, $e^{-\ln 5}$, $e^{3\ln 3}$, $\arctan(\sqrt{3})$, or $\cosh(\ln 3)$ should be simplified.
- \bullet Please show all of your work and justify all of your answers. (You may use the backs of pages for additional work space.)
- **1.** [18 Points] Evaluate each of the following **limits**. Please justify your answer. Be clear if the limit equals a value, $+\infty$ or $-\infty$, or Does Not Exist. Simplify.
- (a) $\lim_{x\to 0} \frac{xe^x \arctan x}{\ln(1+5x) 5x}$ (b) Compute $\lim_{x\to 0} \frac{xe^x \arctan x}{\ln(1+5x) 5x}$ again using series.
- (c) $\lim_{x \to \infty} \left(\frac{x+1}{x}\right)^x$
- 2. [22 Points] Evaluate each of the following integrals.
- (a) $\int \frac{\cos x}{(4+\sin^2 x)^{\frac{5}{2}}} dx$ (b) $\int \frac{x^2}{\sqrt{4-x^2}} dx$ (c) $\int_1^{\sqrt{3}} \frac{1}{\sqrt{4-x^2}} dx$
- **3.** [40 Points] For each of the following **improper integrals**, determine whether it converges or diverges. If it converges, find its value. Simplify.
- (a) $\int_6^7 \frac{8}{x^2 4x 12} dx$ (b) $\int_7^\infty \frac{8}{x^2 4x 12} dx$ Tip: Reuse your algebra work from part (a)
- (c) $\int_0^{e^3} \frac{1}{x \left[3 + (\ln x)^2\right]} dx$ (d) $\int_0^1 \sqrt{x} \ln x dx$
- 4. [18 Points] Find the sum of each of the following series (which do converge). Simplify.
- (a) $\sum_{n=1}^{\infty} \frac{(-1)^n 5^{2n+1}}{2^{5n-1}}$ (b) $\sum_{n=0}^{\infty} \frac{(-1)^{n+1} (\ln 9)^n}{2^{n+1} \cdot n!}$ (c) $\sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n+1}}{9^n (2n)!}$
- (d) $-\frac{1}{3} + \frac{1}{5} \frac{1}{7} + \frac{1}{9} \dots$ (e) $-\frac{\pi^3}{3!} + \frac{\pi^5}{5!} \frac{\pi^7}{7!} + \frac{\pi^9}{9!} \dots$ (f) $-1 + \frac{1}{2} \frac{1}{3} + \frac{1}{4} \frac{1}{5} + \dots$
- 5. [26 Points] In each case determine whether the given series is absolutely convergent, conditionally convergent, or divergent. Justify your answers.
- (a) $\sum_{n=1}^{\infty} (-1)^n \frac{\arctan(7n)}{n^7 + 7}$ (b) $\sum_{n=1}^{\infty} \arctan\left(\frac{n^7 + 1}{n^7 + 7}\right)$ (c) $\sum_{n=1}^{\infty} (-1)^n \left(\frac{n + 1}{n^2}\right)$
- (d) $\sum_{n=1}^{\infty} \frac{(-1)^n (3n)! \ln n}{(n!)^2 e^{4n} n^n}$ (e) $\sum_{n=1}^{\infty} (-1)^n \frac{n^2 + 5}{n^5 + 2}$

- **6.** [18 Points] Find the **Interval** and **Radius** of Convergence for each of the following power series. Analyze carefully and with full justification.
- (a) $\sum_{n=1}^{\infty} \frac{(-1)^n (3x-4)^n}{n^2 \cdot 5^n}$
- (b) $\sum_{n=0}^{\infty} \frac{(-1)^n \ x^{2n}}{(2n)!}$
- (c) $\sum_{n=1}^{\infty} n! (x-6)^n$
- 7. [10 Points] Please analyze with detail and justify carefully. Simplify.
- (a) Use MacLaurin series to **Estimate** $\int_0^1 x \sin(x^2) dx$ with error less than $\frac{1}{1000}$.
- (b) Use MacLaurin Series to **Estimate** $\frac{1}{\sqrt{e}}$ with error less than $\frac{1}{100}$.
- 8. [10 Points] [[] \[\]
- **9.** [18 Points]
- (a) Consider the Parametric Curve represented by $x = \ln t + \ln(1 t^2)$ and $y = \sqrt{8} \arcsin t$.

COMPUTE the **arclength** of this parametric curve for $\frac{1}{4} \le t \le \frac{1}{2}$. Show that the answer simplifies to $\ln\left(\frac{5}{2}\right)$

- 10. [20 Points] For each of the following problems, do the following two things:
- 1. Sketch the Polar curves and shade the described bounded region.
- 2. Set-Up but **DO NOT EVALUATE** the Integral representing the area of the described bounded region.
- (a) The **area** bounded outside the polar curve $r = 1 + \sin \theta$ and inside the polar curve $r = 3 \sin \theta$.
- (b) The **area** bounded outside the polar curve r=2 and inside the polar curve $r=4\sin\theta$.
- (c) The **area** that lies inside both of the curves $r = 1 + \cos \theta$ and $r = 1 \cos \theta$.
- (d) The **area** bounded inside the polar curve $r = 2 + 2\cos\theta$.