

- Please see the course webpage for the answer key.

1. Compute $\int_e^{e^3} \frac{4}{x(\ln x)^2} dx = 4 \int_1^3 \frac{1}{u^2} du = 4 \int_1^3 u^{-2} du = -4u^{-1} \Big|_1^3 = -\frac{4}{u} \Big|_1^3 = -\frac{4}{3} - (-4) = -\frac{4}{3} + 4 = -\frac{4}{3} + \frac{12}{3} = \boxed{\frac{8}{3}}$

Here
$$\begin{array}{l} u = \ln x \\ du = \frac{1}{x} dx \end{array}$$
 and
$$\begin{array}{l} x = e \implies u = \ln e = 1 \\ x = e^3 \implies u = \ln e^3 = 3 \end{array}$$

2. Compute $\int_{\ln 3}^{\ln 8} \frac{e^x}{\sqrt{1+e^x}} dx = \int_4^9 \frac{1}{\sqrt{u}} du = \int_4^9 u^{-\frac{1}{2}} du = 2\sqrt{u} \Big|_4^9 = 2\sqrt{9} - 2\sqrt{4} = 2(3) - 2(2) = 6 - 4 = \boxed{2}$

Here
$$\begin{array}{l} u = 1 + e^x \\ du = e^x dx \end{array}$$
 and
$$\begin{array}{l} x = \ln 3 \implies u = 1 + e^{\ln 3} = 1 + 3 = 4 \\ x = \ln 8 \implies u = 1 + e^{\ln 8} = 1 + 8 = 9 \end{array}$$

3. Compute $\int_{\ln 2}^{\ln 3} \frac{1}{e^{2x} (1 - e^{-2x})^2} dx = \frac{1}{2} \int_{\frac{3}{4}}^{\frac{8}{9}} \frac{1}{u^2} du = -\frac{1}{2u} \Big|_{\frac{3}{4}}^{\frac{8}{9}} = -\frac{1}{2} \left(\frac{1}{(\frac{8}{9})} - \frac{1}{(\frac{3}{4})} \right) = -\frac{1}{2} \left(\frac{9}{8} - \frac{4}{3} \right) = -\frac{1}{2} \left(\frac{27}{24} - \frac{32}{24} \right) = -\frac{1}{2} \left(-\frac{5}{24} \right) = \boxed{\frac{5}{48}}$

Here
$$\begin{array}{l} u = 1 - e^{-2x} \\ du = 2e^{-2x} dx \\ \frac{1}{2}du = \frac{1}{e^{2x}} dx \end{array}$$
 and
$$\begin{array}{l} x = \ln 2 \implies u = 1 - e^{-2 \ln 2} = 1 - e^{\ln(2^{-2})} = 1 - \frac{1}{4} = \frac{3}{4} \\ x = \ln 3 \implies u = 1 - e^{-2 \ln 3} = 1 - e^{\ln(3^{-2})} = 1 - \frac{1}{9} = \frac{8}{9} \end{array}$$

4. Compute $\int \frac{x}{(3x+1)^2} dx = \frac{1}{3} \int \frac{\left(\frac{u-1}{3}\right)}{u^2} du = \frac{1}{9} \int \frac{u-1}{u^2} du = \frac{1}{9} \int \frac{u}{u^2} - \frac{1}{u^2} du = \frac{1}{9} \int \frac{1}{u} - u^{-2} du = \frac{1}{9} \left(\ln |u| + \frac{1}{u} \right) + C = \boxed{\frac{1}{9} \left(\ln |3x+1| + \frac{1}{3x+1} \right) + C}$

$$u = 3x + 1 \Rightarrow x = \frac{u - 1}{3}$$

Here $du = 3dx$

$$\frac{1}{3}du = dx$$

$$\begin{aligned} 5. \quad & \text{Compute } \int_{\frac{\pi}{18}}^{\frac{\pi}{9}} \tan(3x) dx = \int_{\frac{\pi}{18}}^{\frac{\pi}{9}} \frac{\sin(3x)}{\cos(3x)} dx = -\frac{1}{3} \int_{\frac{\sqrt{3}}{2}}^{\frac{1}{2}} \frac{1}{u} du = -\frac{1}{3} \ln|u| \Big|_{\frac{\sqrt{3}}{2}}^{\frac{1}{2}} \\ & = -\frac{1}{3} \left(\ln\left(\frac{1}{2}\right) - \ln\left(\frac{\sqrt{3}}{2}\right) \right) = -\frac{1}{3} \left(\ln\left(\frac{\frac{1}{2}}{\frac{\sqrt{3}}{2}}\right) \right) \\ & = -\frac{1}{3} \left(\ln\left(\frac{1}{\sqrt{3}}\right) \right) = -\frac{1}{3} (\ln 1 - \ln \sqrt{3}) = -\frac{1}{3} (0 - \ln \sqrt{3}) = \boxed{\frac{\ln \sqrt{3}}{3}} \text{ or } \boxed{\frac{\ln 3}{6}} \end{aligned}$$

$$\begin{aligned} 6. \quad & \text{Consider } G(x) = \frac{1}{\sin \sqrt{e^x + e^7}} + \frac{1}{e^{\sqrt{x^2 + 7 \sin x}}} + \frac{1}{\sqrt{7 + e^{\sin x}}} \quad \text{Compute } G'(x). \text{ Do not simplify here.} \\ \text{Here } & \begin{aligned} u &= \cos(3x) \\ du &= -3 \sin(3x) dx \\ -\frac{1}{3}du &= \sin(3x) dx \end{aligned} \quad \text{and} \quad \begin{aligned} x = \frac{\pi}{18} \implies u &= \cos\left(\frac{3\pi}{18}\right) = \cos\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2} \\ x = \frac{\pi}{9} \implies u &= \cos\left(\frac{3\pi}{9}\right) = \cos\left(\frac{\pi}{3}\right) = \frac{1}{2} \end{aligned} \end{aligned}$$

$$6. \quad \text{Consider } G(x) = \frac{1}{\sin \sqrt{e^x + e^7}} + \frac{1}{e^{\sqrt{x^2 + 7 \sin x}}} + \frac{1}{\sqrt{7 + e^{\sin x}}} \quad \text{Compute } G'(x). \text{ Do not simplify here.}$$

First rewrite to simplify and prepare:

$$\begin{aligned} G(x) &= \left(\sin \sqrt{e^x + e^7} \right)^{-1} + e^{-\sqrt{x^2 + 7 \sin x}} + (7 + e^{\sin x})^{-\frac{1}{2}} \\ G'(x) &= \boxed{- \left(\sin \sqrt{e^x + e^7} \right)^{-2} \cos \sqrt{e^x + e^7} \left(\frac{1}{2\sqrt{e^x + e^7}} \right) (e^x)} \text{ (continued ...)} \\ &\quad \boxed{+ e^{-\sqrt{x^2 + 7 \sin x}} \left(-\frac{1}{2\sqrt{x^2 + 7 \sin x}} \right) (2x + 7 \cos x)} \text{ (continued ...)} \\ &\quad \boxed{- \frac{1}{2} (7 + e^{\sin x})^{-\frac{3}{2}} (e^{\sin x}) (\cos x)} \end{aligned}$$

$$7. \quad \text{Consider } F(x) = \sin(\ln(1 + x)) - \frac{1}{1 + \ln(1 + 3x)} \quad \text{Compute the equation of the tangent line to the curve } F(x) \text{ at the point where } x = 0.$$

First compute the corresponding y -value.

$$F(0) = \sin(\ln(1 + 0)) - \frac{1}{1 + \ln(1 + 0)} = \sin(0) - \frac{1}{1 + 0} = 0 - 1 = -1$$

Second, compute the derivative for $F(x) = \sin(\ln(1 + x)) - (1 + \ln(1 + 3x))^{-1}$

$$F'(x) = \cos(\ln(1+x)) \left(\frac{1}{1+x} \right) + \frac{1}{(1+\ln(1+3x))^2} \left(\frac{1}{1+3x} \right) \quad (3)$$

Next find the specific slope when $x = 0$.

$$\begin{aligned} F'(0) &= \cos(\ln(1+0)) \left(\frac{1}{1+0} \right) + \frac{1}{(1+\ln(1+0))^2} \left(\frac{1}{1+0} \right) \quad (3) \\ &= \cos(0)(1) + \frac{1}{(1+0)^2}(1)(3) = (1)(1) + (1)(1)(3) = 4 \end{aligned}$$

Finally, use *point-slope form* to get the equation of the tangent line $y - (-1) = 4(x - 0)$.

We have $\boxed{y = 4x - 1}$